

## *Discussion*



## Discussion

Although asthma prevalence data in North Carolina are demonstrating downward trends, asthma is still a huge issue in the state. In 2005, 6.5% of adults and 11.5% of children reported currently having asthma. While the percentage of adults with current asthma in North Carolina is below the national median, the percentage of children in North Carolina with asthma exceeds the national median number from the most recent national numbers (2004).

Although asthma affects the whole population, certain subgroups appear to be more adversely affected than others. Gender differences are seen in each of the surveillance measures in this document. Male children in North Carolina have a higher asthma prevalence than female children. Conversely, adult females have a higher prevalence of asthma than adult men in North Carolina. Females also have a higher rate of hospitalization due to a primary cause of asthma than men, as well as a higher rate of mortality due to asthma than North Carolina men.

Age is also factor when looking at who asthma affects. The very young, ages 0 to 4, have the highest rates of hospitalization due to a primary cause of asthma. Adults age 65 and older have the second highest asthma hospitalization rate, following the 0 to 4 age group. Adults in the age group 65+ have a significantly higher mortality due to a primary cause of asthma than all other age groups in North Carolina.

Disparities between racial groups, especially between African Americans, Native Americans and whites in North Carolina, are an issue. According to the 2005 N.C. YRBS, both male and female African American high school students have higher prevalence of asthma than white high school students. Although the 2005 N.C. BRFSS data did not show a statistically significant difference for Native Americans having a higher

asthma prevalence of asthma than whites, it has been shown in previous years. This data is consistent with national estimates that show Native Americans are 25% more likely to have been diagnosed with asthma than whites.<sup>12</sup>

Exploring health care utilization data, according to the 2005 N.C. BRFSS, African American adults were more likely than white adults to visit an emergency room or urgent care center three or more times in the previous year because of their asthma. African American children were also more likely than white children to have visited an emergency room or urgent care center in the past year because of asthma. Unfortunately, due to inconsistency in reporting race and ethnicity data, we cannot provide racial and ethnic data on actual hospitalizations.

Although there were only 116 deaths due to asthma in 2005 in North Carolina, African Americans were disproportionately affected. The mortality rate due to a primary cause of asthma in 2005 for African Americans was more than twice that of whites. Looking at the past 6 years (1999-2005), African Americans<sup>a</sup> had a mortality rate due to asthma 2.5 times higher than whites. Minorities<sup>b</sup> have a higher mortality due to a primary cause of asthma in each age group of North Carolinians age 5 and older.

Additionally, low income households were disproportionately affected by asthma. North Carolina adults with a household income less than \$15,000 a year were more likely to have higher asthma prevalence than all other income groups. This is similar to national trends, which show that adults in poor families have higher percentages of asthma than adults in families that are not poor.<sup>13</sup>

Currently, very little data is available on provider practices and asthma management, specifically providing each child or adult with asthma with an up to date asthma management plan. According to the 2005 N.C. CHAMP survey, 43% of children

<sup>a</sup>This rate includes Asians and other minorities. These groups make up for such a small number of deaths over the 6 year (23) that they were grouped with African Americans.

<sup>b</sup>The minority group being discussed here is comprised mainly of African Americans (328), but also includes Asians, American Indians, and other non-white racial groups.

with current asthma in North Carolina have reported (parental/guardian report) not receiving an asthma management plan from their physician or other health care professional. North Carolina Asthma Program is working on obtaining similar information from adults, and questions will be included on future statewide telephone surveys, as well in other physician specific project that the Chronic Disease and Injury Section of the North Carolina Division of Public Health, is conducting.

## ***Limitations***

While the *Burden of Asthma in North Carolina* is a comprehensive report, there are data gaps and limitations. Data gaps include limited available data on Native Americans, lack of prevalence data on asthma triggers (outside of tobacco smoke), lack of information on health care providers' practices, and a lack of accurate race and ethnicity data in hospitalization records.

Other data gaps concerning hospitalization records include the inability to track repeat visits for persons with asthma to the emergency room or admissions to the hospital. The ability to identify who with current asthma is making multiple emergency room visits and hospitalized repeatedly, would be an extremely beneficial tool for determining asthma management techniques. There is currently no way to track how many repeat visits a patient makes other than self report through the N.C. BRFSS.

Lack of data regarding work related asthma is also an issue in North Carolina. Several methods are currently being investigated as possible methods for filling this gap. The North Carolina Asthma Program will look at how other states are satisfying this data need and try to determine if any of those methods would be appropriate here.

## ***Next Steps***

The North Carolina Asthma Program will work together with the statewide coalition, the North Carolina Asthma Alliance, and other partners and stakeholders, to use this data to develop a state asthma plan. The North Carolina State Asthma Plan, that will be available March 2007, will address strategies for reducing the burden of asthma in North Carolina, with a specific emphasis on the disproportionately affected groups described herein. The State Asthma Plan will also provide methods for addressing the limitations discussed above.

The North Carolina Asthma Program will continue to conduct surveillance across the state utilizing a variety of available methods. We will work to fill any data gaps, while continuing to make the most out of the data that we already have. The Asthma Program will also investigate new ways to capture populations that we know are being negatively affected by asthma, but we do not have sufficient data on; including Native Americans and low income households.



## *References*



## References

1. Boss, L.P., Kreutzer, R.A., Luttinger, D., Leighton, J., Wilcox, K., Redd, S.C. (2001) The Public Health Surveillance of Asthma. *Journal of Asthma*, 38(1), 83-89.
2. Gordis, L. (2000) *Epidemiology*. 2nd ed. Philadelphia, Pennsylvania: W.B. Saunders Company.
3. CSTE POSITION STATEMENT 1998-EH/CD 1: Asthma Surveillance and Case Definition. Retrieved 5/24/2006. From Council for State and Territorial Epidemiologists. Web page. <http://www.cste.org/ps/1998/1998-eh-cd-01.htm>.
4. Centers for Disease Control and Prevention (CDC). 2004 National asthma training curriculum. Version 1. [CD ROM]. Atlanta; CDC; 2004.
5. North Carolina State Center for Health Statistics (SCHS). 2005 Behavioral Risk Factor Surveillance System (BRFSS). Retrieved 6/5/2006. Web Page: <http://www.schs.state.nc.us/schs/brfss/>.
6. North Carolina State Center for Health Statistics (SCHS). 2004 BRFSS Annual Results Technical Notes. Retrieved 6/5/2006. Web Page: <http://www.schs.state.nc.us/SCHS/brfss/2004/technical.html>.
7. Rhodes, L., Moorman, J.E., Redd, S.C., Mannino, D.M. (2003) Self-Reported Asthma Prevalence and Control Among Adults—United States, 2001. *MMWR*, 52 (17), 381-384.
8. Asthma in Minnesota: 2005 Epidemiology Report. Minnesota Department of Health. Minneapolis, MN. September 2005.
9. National Institute of Health (NIH) (1999) *National Heart, Lung, and Blood Institute Data Fact Sheet: Asthma Statistics*. Retrieved 6/5/2006. From National Heart, Lung, and Blood Institute. Web site: <http://www.nhlbi.nih.gov/health/prof/lung/asthma/asthstat.pdf>.
10. Centers for Disease Control and Prevention (CDC). 2005 *Behavioral Risk Factor Surveillance System Survey Data*. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, 2005.
11. State Center for Health Statistics (SCHS) and Office of Minority Health and Health Disparities. (OMH) 2005 *North Carolina Minority Health Facts: American Indians*. North Carolina Department of Health and Human Services. Retrieved 6/5/2006. Web site: <http://www.schs.state.nc.us/SCHS/pdf/FactsAI2005.pdf>.
12. National Center for Health Statistics (NCHS). (2002) *Asthma Prevalence, Health Care Use and Mortality, 2002*. Centers for Disease Control and Prevention. Retrieved 6/30/2006. Web site: <http://www.cdc.gov/nchs/products/pubs/pubd/hestats/asthma/asthma.htm>.
13. Lethbridge-Cejku M, Rose D, Vickerie J. (2006) *Summary health statistics for U.S. Adults: National Health Interview Survey, 2004*. National Center for Health Statistics. Vital Health Stat 10(229).
14. Bloom B, Dey AN. (2006) *Summary Health Statistics for U.S. Children: National Health Interview Survey, 2004*. National Center for Health Statistics. Vital Health Stat. 10(227).

15. Centers for Disease Control and Prevention (CDC). (2006) *Youth Risk Behavior Surveillance – United States, 2005*. Surveillance Summaries, June 9, 2006. MMWR 2006; 55 (No. SS-5).
16. National Heart, Lung, and Blood Institute (NHLBI). 1997. Expert Panel Report 2: Guidelines for the Diagnosis and Management of Asthma. National Institutes of Health, Bethesda, MD. Web site: <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.pdf>.
17. Asthma and Allergy Foundation of America (AAFA), Florida Chapter, Inc. *Adult Onset Asthma*. Retrieved 5/25/2006. Web site: [http://www.aafaflorida.org/features/answers/adult\\_onset\\_asthma.htm](http://www.aafaflorida.org/features/answers/adult_onset_asthma.htm).
18. National Heart, Lung, and Blood Institute (NHLBI). (2003). *Guidelines for the Diagnosis and Management of Asthma, Update on Selected Topics, 2002*. National Institutes of Health, Bethesda, MD. Retrieved 7/10/2006. Web site: <http://www.nhlbi.nih.gov/guidelines/asthma/asthmafullrpt.pdf>.
19. National Heart, Lung, and Blood Institute. 1997. *Problem Sleepiness in Your Patient*. National Institutes of Health, Bethesda, MD.
20. Janson C, Gislason T, Boman G, Hetta J, Roos BE. Sleep disturbances in patients with asthma. *Respiratory Medicine*. 1990 Jan/ 84 (1): 37-42.
21. Centers for Disease Control and Prevention. *Measuring Healthy Days*. Atlanta, Georgia: CDC, November 2000.
22. Ford, ES, Mannino, DM, Homa, DM, Gwynn, C, Redd, SC, Moriarty, DB, Mokdad, AH. Self-reported asthma and health-related quality of life: Findings from the behavioral risk factor surveillance system. *Chest* 2003; 123 (1): 119-127.
23. N.C. Department of Health and Human Resources, Division of Public Health. *The Health of North Carolinians: A Profile*. Retrieved 5/16/2006. Web Page: <http://www.schs.state.nc.us/SCHS/pdf/HealthProfile.pdf>.
24. Asthma Initiative of Michigan. *For Healthy Lungs: Written Asthma Actions Plan Components and Distribution*. Retrieved 7/12/2006. Web page: [http://www.getastmahelp.org/actionplan\\_components.asp](http://www.getastmahelp.org/actionplan_components.asp).
25. Asthma and Allergy Foundation of America. *Asthma Overview*. Retrieved 6/17/2006. Web site: <http://aafa.org/>.
26. Centers for Disease Control and Prevention. *Strategies for Addressing Asthma Within a Coordinated School Health Program with Updated Resources*. Atlanta, Georgia: Centers for Disease Control and Prevention. National Center for Chronic Disease Prevention and Health Promotion, 2005. Available at: [www.cdc.gov/healthyYouth/asthma/pdf/strategies.pdf](http://www.cdc.gov/healthyYouth/asthma/pdf/strategies.pdf).
27. Public Schools of North Carolina, Department of Public Instruction. *North Carolina's School Health Education Profile: 2004 Principal's Survey*. Department of Health and Human Services, Division of Public Health.
28. National Heart, Lung, and Blood Institute, National Institutes of Health. *Asthma and Physical Activity in the School: Making a Difference*. NIH Publication No. 95-3651. September 1995.

29. Environmental Protection Agency. *Asthma and Outdoor Air Pollution*. Retrieved 5/31/2006. Web page: <http://www.epa.gov/asthma/outdoorair.html>.
30. Environmental Protection Agency. Indoor Environmental Asthma Triggers. Retrieved 5/31/2006. Web page: <http://www.epa.gov/asthma/triggers.html>.
31. NC Department of Health and Human Services and the University of North Carolina School of Public Health. *Asthma in North Carolina: The North Carolina School Asthma Survey, 1999-2000*. Retrieved 5/17/2006. Web site: <http://wch.dhhs.state.nc.us/Asthma/pdf/Leg%204%20Pager.pdf>.
32. Environmental Protection Agency. Indoor Environmental Asthma Triggers – Secondhand Smoke. Retrieved 5/31/2006. Web page: <http://www.epa.gov/asthma/shs.html>.
33. U.S. Department of Health and Human Services. The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2006.
34. U.S. Department of Health and Human Services. *The Health Consequences of Smoking: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2004.
35. U.S. Department of Health and Human Services, National Heart, Lung, and Blood Institute. U.S. Department of Education. Revised 2003. National Asthma Education and Prevention Program: Managing Asthma – A Guide For Schools. Available at: [http://www.nhlbi.nih.gov/health/prof/lung/asthma/asth\\_sch.pdf](http://www.nhlbi.nih.gov/health/prof/lung/asthma/asth_sch.pdf).
36. Centers for Disease Control and Prevention. Tobacco Information and Prevention Source. Retrieved 8/10/2006. Web Page: [http://www.cdc.gov/tobacco/overview/Fast\\_Facts.htm](http://www.cdc.gov/tobacco/overview/Fast_Facts.htm).
37. U.S. Department of Health and Human Services, Health Resources and Services Administration, Maternal and Child Health Bureau. The National Survey of Children's Health 2003. Rockville, Maryland: U.S. Department of Health and Human Services, 2005.
38. North Carolina Tobacco Prevention and Control Branch. *North Carolina 2005 Youth Tobacco Survey (YTS): High School Fact Sheet*. Division of Public Health, Department of Health and Human Services.
39. North Carolina Tobacco Prevention and Control Branch. *North Carolina 2005 Youth Tobacco Survey (YTS): Middle School Fact Sheet*. Division of Public Health, Department of Health and Human Services.
40. Weiss KB, Gergen, PJ, Hodgson T. An economic evaluation of asthma in the United States. *New England Journal of Medicine* 1992; 326: 862-866.
41. Akinbami, L.J., Schoendorf, K.C. *Trends in Childhood Asthma: Prevalence, Health Care Utilization, and Mortality*. *Pediatrics* 2002; 110: 315-322.



42. Coffey, RM, Ho, K, Adamson, DM, Matthews, TL, Sewell, J. *Asthma Care Quality Improvement: A Resource Guide for State Action*. (Prepared by Thomson Medstat and The Council of State Governments under Contract No. 290-000-0004). Rockville, MD: Agency for Healthcare Research and Quality, Department of Health and Human Services; April 2006. AHRQ Pub No. 06-0012-1.
43. Grunbaum JA, Di Pietra J, McManus T, Hawkins J, Kann L. *School Health Profiles: Characteristics of Health Programs Among Secondary Schools (Profiles 2004)*. Atlanta, GA: Centers for Disease Control and Prevention, 2005.
44. North Carolina Emergency Department Database. Retrieved 7/12/2006. Web page: <http://www.ncedd.org/>.
45. National Committee for Quality Assurance. Retrieved 7/17/2006 Web site: <http://www.ncqa.org/programs/hedis/>.
46. National Committee for Quality Assurance (NCQA). HEDIS 2004. Health plan employer data & information set. Vol. 2, Technical specifications. Washington (DC): National Committee for Quality Assurance (NCQA); 126 p.
47. South Carolina Department of Health and Environmental Control. South Carolina Community Assessment Network. Death Certificate Definitions. Retrieved July 19, 2006. Web Page: <http://scangis.dhec.sc.gov/scanet/defn/dthdefn.htm#mult%20cause>.
48. U.S. Department of Health and Human Services. *Healthy People 2010: Understanding and Improving Health*. 2nd ed. Washington, DC: U.S. Government Printing Office, November 2000.
49. Healthy Carolinians. North Carolina 2010 Health Objectives. North Carolina Division of Public Health. Retrieved 7/20/2006. Web site: <http://www.healthycarolinians.org/healthobj2010.htm>.
50. Centers for Disease Control and Prevention (CDC). *Behavioral Risk Factor Surveillance System Survey Data*. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, 2000-2005.
51. Herrick, H, Gizlice, Z. North Carolina State Center For Health Statistics. *Spanish-Speaking Hispanics in North Carolina: Health Status, Access to Health Care, and Quality of Life*. Results from the 2002 and 2003 NC BRFSS Surveys. North Carolina Division of Public Health. Retrieved 8/18/2006. Web site: <http://www.schs.state.nc.us/SCHS/pdf/SCHS143.pdf>.
52. State Center for Health Statistics. *Behavioral Risk Factor Surveillance System (BRFSS) 2005 Results*. North Carolina Department of Health and Human Services, Division of Public Health. Retrieved 8/18/2006. Web site: <http://www.schs.state.nc.us/SCHS/brfss/2005/index.html>.
53. State Center for Health Statistics. *Child Health Assessment and Monitoring Program (CHAMP) 2005 Results*. North Carolina Department of Health and Human Services, Division of Public Health. Retrieved 8/18/2006. Web site: <http://www.schs.state.nc.us/SCHS/champ/2005/topics.html>.
54. Department of Public Instruction. *Youth Risk Behavior Survey (YRBS) 2005*. North Carolina Department of Health and Human Services. Retrieved 6/27/2006. Web site: <http://www.nchealthyschools.org/data/yrbs/>.

55. Yeatts, K, Shy, C, Sotir, M, Music, S, Herget, C. *Health Consequences for Children with Undiagnosed Asthma-Like Symptoms*. Archives of Pediatric Adolescent Medicine 2003; 157: 540-544.
56. American Lung Association. *Trends in Asthma Morbidity and Mortality*. Epidemiology and Statistics Unit, Research and Program Services. July 2006. Retrieved 9/25/2006. Web site: <http://www.lungusa.org/atf/cf/%7B7A8D42C2-FCCA-4604-8ADE-7F5D5E762256%7D/ASTHMA06FINAL.PDF>.
57. Centers for Disease Control and Prevention. *Asthma: Basic Facts*. Environmental Hazards and Health Effects Program, Air Pollution and Respiratory Health Branch. Retrieved 6/5/2006. Web site: <http://www.cdc.gov/asthma/faqs.htm>.
58. National Heart, Lung, and Blood Institute (NHLBI). *Disease and Conditions Index: Asthma*. National Institutes of Health, Bethesda, MD. Retrieved 6/15/2006. Web site: [http://www.nhlbi.nih.gov/health/dci/Diseases/Asthma/Asthma\\_WhatIs.html](http://www.nhlbi.nih.gov/health/dci/Diseases/Asthma/Asthma_WhatIs.html).
59. Interview with Dr. Gerri Mattson, M.D, MSPH, Pediatric Medical Consultant, Children and Youth Branch, NC Division of Public Health. April, 2006.
60. National Toxicology Program. 9th Report on Carcinogens, 2000. Research Triangle Park, NC: U.S. Department of Health and Human Sciences, National Institute of Environmental Health Sciences, 2000. Retrieved June 2006. Web site: <http://ntp.niehs.nih.gov/ntp/roc/elev-enth/profiles/s176toba.pdf>

## *Technical Notes*



## Technical Notes

### Part 1. Odds Ratio

In a study where participants are selected on the basis of their disease status, as in the N.C. BRFSS, the relative risk can be estimated by calculating the ratio of the odds of exposure among the cases to that among the controls. In this document, cases are considered persons who have either lifetime or current asthma, and exposure is gender.<sup>1</sup>

Example – based on the whether a female has a greater odds of having asthma than a male.

	Has Asthma	Does Not Have Asthma
Female	<b>A</b>	<b>B</b>
Male	<b>C</b>	<b>D</b>

Odds Ratio (OR) =  $\frac{\text{Odds that an exposed person (female) develops the disease (asthma)}}{\text{Odds that a non-exposed person (male) develops the disease (asthma)}}$   
 $OR = A \cdot D / B \cdot C$

### Part 2. Prevalence

Prevalence is defined as the number of affected persons present in the population at a specific time divided by the number of persons in the population at that time. It is used to describe the health burden on a specific population.<sup>2</sup>

Prevalence =  $\frac{\text{\# of cases of a disease present in the population at a specified time}}{\text{\# of persons in the population at that specified time}}$

### Part 3. Age Adjustment

Populations often differ in age distribution. Therefore, it is often important to control for the differences among the age distributions of populations when making comparisons among death rates to assess the relative risk of death. The direct method of age-adjustment is frequently used to compare the death rates of different populations, by controlling for differences in age distribution. Sum the products of the age-specific death rates and the proportion of the standard population in that age group across all ten age groups. This weighted sum is represented in the following formula:

$$\text{Age-adjusted death rate} = \sum_{i=1}^{10} w_i p_i$$

Where:  $p_i$  = the age specific rate for age group  $i$ .

$w_i$  = the weight; the proportion of the standard population in age group  $i$ .<sup>5</sup>

The standard population used to calculate age-adjusted rates in this document is the 2000 United States Standard Population.

### ***Part 4. Confidence Intervals for Proportions***

The confidence interval represents the range within which the true magnitude of effect lies with a certain degree of assurance. A 95% confidence interval states that we are 95% certain that the true measure lies within this specified range.<sup>1</sup>

For example, the estimated current asthma prevalence among North Carolina adults (from a random sample of the population) is 6.5%, with a 95% confidence interval of 6.0% to 7.0%. This means that we are 95% confident that the true prevalence of current asthma for North Carolina adults is no less than 6.0%, and no greater than 7.0%. A 95% confidence uses a multiplier of 1.96.

The formula for the 95% confidence interval is:

$$p \pm 1.96 \sqrt{\frac{pq}{n}}$$

Where: p = proportion

n = sample size

q = 1-p (for small values of p ( $\leq .01$ ) q is small and may be ignored)<sup>4</sup>

### ***Part 5. Confidence Intervals for Death Rates***

Confidence intervals are used when looking at the age adjusted death rates in this document. The formula is the age-adjusted proportion of persons who died in this time period (p) plus/minus 1.96 (for a 95% confidence interval) multiplied by the standard error of an age-adjusted death rate, which is:

$$RSE(R'') = 100 \frac{\sqrt{\sum \left\{ w_i^2 R_i^2 \left( \frac{1}{D_i} \right) \right\}}}{R''}$$

Approximate 95% Confidence Interval: 1-99 deaths

Lower:  $R'' * L(1 - \alpha = .95, D_{adj})$

Upper:  $R'' * U(1 - \alpha = .95, D_{adj})$

where

$R''$  = age-adjusted rate (per 100,000 population) =  $\sum w_i R_i$

$w_i$  =  $i^{th}$  age-specific Standard Population such that  $\sum(w_i) = 1.0$

$R_i$  = age-specific rate (per 100,000) for the  $i^{th}$  age group

$D_i$  = total number of deaths for the  $i^{th}$  age group upon which age-specific rate is based

$$S(R'') = R'' * \frac{RSE(R'')}{100} = \text{standard error of age-adjusted rate}$$

$L(1 - \alpha = .95, D_{adj})$  and  $U(1 - \alpha = .95, D_{adj})$  are lower and upper 95% confidence limit factors and are shown in table S

$$D_{adj} = \frac{1}{\left( \frac{RSE(R'')}{100} \right)^2} = \text{adjusted number of deaths rounded to nearest integer}$$

Table S found in Technical Appendix from the *Vital Statistics of United States 1999 Mortality*.<sup>6</sup>

Age-Specific confidence intervals for less than 100 deaths.

$$RSE(R) = RSE(D) = 100 \sqrt{\frac{1}{D}}$$

Approximate 95% Confidence Interval: 1-99 deaths

Lower:  $R * L(1 - \alpha = .95, D)$

Upper:  $R * U(1 - \alpha = .95, D)$

where

$R$  = rate (deaths per 100,000 population)

$D$  = total number of deaths upon which rate is based

$$S(R) = R * \frac{RSE(R)}{100} = \text{standard error of rate}$$

$L(1 - \alpha = .95, D)$  and  $U(1 - \alpha = .95, D)$  are lower and upper 95% confidence limit factors and are shown in table S

Table S found in Technical Appendix from the *Vital Statistics of United States 1999 Mortality*.<sup>6</sup>

## ***Part 6. Trend Analysis***

The Spearman Rank Order Correlation test was utilized to determine if there was a trend in total mortality rates from 1995 through 2005, and for age, race, and gender specific mortality rates from 1999 through 2005. This test quantifies the extent to which there is a linear relationship between the rate and year.<sup>1</sup>

The correlation coefficient ( $\rho$ ,  $\rho$ ) can vary between +1.0 and -1.0. If the coefficient equals -1.0, it indicates a perfect negative correlation, with each year having a lower mortality rate for that specific group than the previous year, for example. If the coefficient equals +1.0, it indicates a perfect positive correlation, where each year has a higher mortality rate for that specific group than the previous year. As the correlation coefficient approaches 0.0, from either direction, the relationship between the variables weakens.<sup>3</sup>

The p-value for the Spearman Rank Order Correlation test ranges from 0.0 to 1.0, and gives the probability of finding a significant overall trend when no trend actually exists. The standard used to assess the significance of a statistical test is a p-value of 0.05. A p-value less than or equal to 0.05 indicates that there is at most a 5% chance of observing a trend that, in reality, does not exist. If the p value is greater than 0.05, chance cannot be excluded as a likely explanation for the trend, so the result is not statistically significant.<sup>3</sup>

## ***References***

1. Hennekens, C, Buring, J. (1987). *Epidemiology in Medicine*. Boston, MA: Little, Brown, and Company.
2. Gordis, L. (2000) *Epidemiology*. 2nd ed. Philadelphia, Pennsylvania: W.B. Saunders Company.
3. Wasilevich, E, Lyon-Callo, S. *Epidemiology of Asthma in Michigan: 2004 Surveillance Report*. Michigan Department of Community Health. June 2004.
4. Buescher, P. 1997. *Problem With Rates Based On Small Numbers*. State Center for Health Statistics. North Carolina Division of Public Health. North Carolina Department of Health and Human Services. Retrieved 10/4/06. Web site: <http://www.schs.state.nc.us/SCHS/pdf/primer12.pdf>.
5. Buescher, P. 1998. *Age-Adjusted Death Rates*. State Center for Health Statistics. North Carolina Division of Public Health. North Carolina Department of Health and Human Services. Retrieved 10/4/06. Web site: <http://www.schs.state.nc.us/SCHS/pdf/primer13.pdf>.
6. Mortality Statistics Branch. *Technical Appendix from the Vital Statistics of United States 1999 Mortality*. National Center for Health Statistics. Centers for Disease Control and Prevention. Department of Health and Human Services. Retrieved 2/14/2006. Web site: <http://www.cdc.gov/nchs/data/statab/techap99.pdf>.